

Cuntz-Pimsner C^* -algebras associated with subshifts

Toke Meier Carlsen*

*Institut for Matematiske Fag, Københavns Universitet
Universitetsparken 5, 2100 København Ø, Denmark
toke@math.ku.dk*

Abstract

By using C^* -correspondences and Cuntz-Pimsner algebras, we associate to every subshift (also called a shift space) X a C^* -algebra $\mathcal{O}_X$, which is a generalization of the Cuntz-Krieger algebras. We show that $\mathcal{O}_X$ is the universal C^* -algebra generated by partial isometries satisfying relations given by X . We also show that $\mathcal{O}_X$ is a one-sided conjugacy invariant of X .

Keywords: C^* -algebras, subshifts, shift spaces, conjugacy, Cuntz-Krieger algebras, Cuntz-Pimsner algebras.

Mathematics Subject Classification 2000: Primary: 46L55, Secondary: 37B10.

1 Introduction

In [7] Cuntz and Krieger introduced a new class of C^* -algebras which in a natural way can be viewed as universal C^* -algebras associated with subshifts (also called shift spaces) of finite type. From the point of view of operator algebra these C^* -algebras were important examples of C^* -algebras with new properties and from the point of view of topological dynamics these C^* -algebras (or rather, the K -theory of these C^* -algebras) gave new invariants of subshifts of finite type.

*Current address: Mathematisches Institut, Einsteinstraße 62, 48149 Münster, Germany

In [13] Matsumoto tried to generalize this idea by constructing C^* -algebras associated with every subshift, and he and others have studied these C^* -algebras in [2, 10, 14–21]. Unfortunately there is a mistake in [17] which makes many of the results in [16–19, 21] invalid for the C^* -algebras constructed in [13]. This mistake has to do with the identification of an underlying compact space which among other things determine the K -theory of the C^* -algebras. It turned out that this compact space is not the space Matsumoto thought it was, and thus many of the results of [16–19, 21] are invalid for the C^* -algebras constructed in [13]. To recover these results Matsumoto and the author introduced in [5] a new class of C^* -algebras associated with subshifts, which has the right underlying compact space and thus satisfies most of the results in [2, 10, 13–21], but these C^* -algebras do not in general have the universal property (cf. Theorem 7.2, Remark 7.3 and 7.4, [13, Theorem 4.9] and [5, pp. 148–149]). Thus it is natural to think of this class of C^* -algebras as the class of reduced C^* -algebras associated with subshifts.

In this paper we will for each subshift X construct a new C^* -algebra $\mathcal{O}_X$ by using C^* -correspondences (also called Hilbert bimodules) and Cuntz-Pimsner algebras, and this new C^* -algebra will both have the right underlying compact space and have the universal property and hence will satisfy all the results of [2, 10, 13–21] and has the C^* -algebra defined in [5] as a quotient. Thus it seems right to think of this C^* -algebra as the universal C^* -algebra associated to a subshift.

The C^* -algebra $\mathcal{O}_X$ can also be constructed as the C^* -algebra of a groupoid (cf. Remark 7.7), and by using Exel's crossed product of a C^* -algebra of an endomorphism (cf. Remark 7.6).

Matsumoto's original construction associated a C^* -algebra to every *two-sided* subshift, but it seems more natural to work with *one-sided* subshifts, so we will do that in this paper, but since every two-sided subshift comes with a canonical one-sided subshift (see below), the C^* -algebras we define in this paper can in a natural way also be seen as C^* -algebras associated to two-sided subshifts (cf. Remark 7.4).

We will show that $\mathcal{O}_X$ is the universal C^* -algebra associated with partial isometries satisfying relations giving by X and which resemble the Cuntz-Krieger relations (Theorem 7.2). We will also show that $\mathcal{O}_X$ is an invariant of X in the sense that if X and Y are conjugate one-sided subshifts, then $\mathcal{O}_X$ and $\mathcal{O}_Y$ are isomorphic (Theorem 8.6). This is a generalization of [7, Proposition 2.17] and [13, Proposition 5.8] (see [16, Lemma 4.5] for a proof of the later Proposition), where it is required that X and Y satisfy a certain condition (I).

2 Notation and preliminaries

Throughout this paper, $\mathbb{N}_0$ will denote the set of non-negative integers.

Let $\mathfrak{a}$ be a finite set endowed with the discrete topology. We will call this set the alphabet. Let $\mathfrak{a}^{\mathbb{N}_0}$ be the infinite product spaces $\prod_{n=0}^{\infty} \mathfrak{a}$ endowed with the product topology. The transformation σ on $\mathfrak{a}^{\mathbb{N}_0}$ given by $(\sigma(x))_i = x_{i+1}$, $i \in \mathbb{N}_0$ is called the *shift*. Let X be a shift invariant closed subset of $\mathfrak{a}^{\mathbb{N}_0}$ (by shift invariant we mean that $\sigma(X) \subseteq X$, not necessarily $\sigma(X) = X$). The topological dynamical system $(X, \sigma|_X)$ is called a *subshift*. We will denote $\sigma|_X$ by σ_X or σ for simplicity, and on occasion the alphabet $\mathfrak{a}$ by $\mathfrak{a}_X$. Since σ_X maps X into X , we can compose σ_X with itself. We will denote this map by σ_X^2 and in general for any positive integer n the n -fold composition of σ_X with itself by σ_X^n . We will for a subset Y of the subshift X and an integer n by $\sigma^n(Y)$ denote

$$\sigma^n(Y) = \begin{cases} \sigma_X^n(Y) & n > 0, \\ Y & n = 0, \\ (\sigma_X^{-n})^{-1}(Y) & n < 0. \end{cases}$$

A finite sequence $\mu = (\mu_1, \dots, \mu_k)$ of elements $\mu_i \in \mathfrak{a}$ is called a finite *word*. The *length* of μ is k and is denoted by $|\mu|$. We let for each $k \in \mathbb{N}_0$, $\mathfrak{a}^k$ be the set of all words with length k and we let $L^k(X)$ be the set of all words with length k appearing in some $x \in X$. We set $L_l(X) = \bigcup_{k=0}^l L^k(X)$ and $L(X) = \bigcup_{k=0}^{\infty} L^k(X)$ and likewise $\mathfrak{a}_l = \bigcup_{k=0}^l \mathfrak{a}^k$ and $\mathfrak{a}^* = \bigcup_{k=0}^{\infty} \mathfrak{a}^k$, where $L^0(X) = \mathfrak{a}^0$ denote the set consisting of the empty word ϵ which has length 0. $L(X)$ is called the *language* of X . Note that $L(X) \subseteq \mathfrak{a}^*$ for every subshift.

For a subshift X and a word $\mu \in L(X)$ we denote by $C_X(\mu)$ the *cylinder set*

$$C_X(\mu) = \{x \in X \mid (x_1, x_2, \dots, x_{|\mu|}) = \mu\}.$$

It is easy to see that

$$\{C_X(\mu) \mid \mu \in L(X)\}$$

is a basis for the topology of X , and that $C_X(\mu)$ is closed and hence compact for every $\mu \in L(X)$. We will allow us self to write $C(\mu)$ instead of $C_X(\mu)$ when it is clear which subshift space we are working with.

For a subshift X and words $\mu, \nu \in L(X)$ we denote by $C(\mu, \nu)$ the set

$$C(\nu) \cap \sigma^{-|\nu|}(\sigma^{|\mu|}(C(\mu))) = \{\nu x \in X \mid \mu x \in X\}.$$

If X and Y are two subshifts and $\phi : X \rightarrow Y$ is a homeomorphism such that $\psi \circ \sigma_X = \sigma_Y \circ \phi$, then we say that ϕ is a *conjugacy* and that X and Y are *conjugate*.

What we have defined above is a *one-sided* subshift. A *two-sided* subshift is defined in the same way, except that we replace $\mathbb{N}_0$ with $\mathbb{Z}$: Let $\mathfrak{a}^{\mathbb{Z}}$ be the infinite product spaces $\prod_{n=-\infty}^{\infty} \mathfrak{a}$ endowed with the product topology, and let σ be the transformation on $\mathfrak{a}^{\mathbb{Z}}$ given by $(\sigma(x))_i = x_{i+1}$, $i \in \mathbb{Z}$. A shift invariant closed subset Λ of $\mathfrak{a}^{\mathbb{Z}}$ (here, by shift invariant we mean $\sigma(\Lambda) = \Lambda$) is called a *two-sided subshift*. The set

$$X_{\Lambda} = \{(x_i)_{i \in \mathbb{N}_0} \mid (x_i)_{i \in \mathbb{Z}} \in \Lambda\}$$

is a one-sided subshift, and it is called the one-sided subshift of Λ .

3 Cuntz-Pimsner algebras

We will in this section give a short introduction to Cuntz-Pimsner algebras. We will follow the universal approach of [8] (see also [23], [24] and [11]).

Let $\mathcal{A}$ be a C^* -algebra. A right *Hilbert $\mathcal{A}$ -module* H is a Banach space with a right action of the C^* -algebra $\mathcal{A}$ and an $\mathcal{A}$ -valued inner product $\langle \cdot, \cdot \rangle$ satisfying

1. $\langle \xi, \eta a \rangle = \langle \xi, \eta \rangle a$,
2. $\langle \xi, \eta \rangle = \langle \eta, \xi \rangle^*$,
3. $\langle \xi, \xi \rangle \geq 0$ and $\|\xi\| = \|\langle \xi, \xi \rangle\|^{1/2}$,

for $\xi, \eta \in H$ and $a \in \mathcal{A}$.

For a Hilbert $\mathcal{A}$ -module H , we denote by $\mathcal{L}(H)$ the C^* -algebra of all adjointable operators on H . For $\xi, \eta \in H$, the operator $\theta_{\xi, \eta} \in \mathcal{L}(H)$ is defined by $\theta_{\xi, \eta}(\zeta) = \xi \langle \eta, \zeta \rangle$ for $\zeta \in H$. We define $\mathcal{K}(H) \subseteq \mathcal{L}(H)$ by

$$\mathcal{K}(H) = \overline{\text{span}}\{\theta_{\xi, \eta} \mid \xi, \eta \in H\},$$

where $\overline{\text{span}}\{\dots\}$ means the closure of the linear span of $\{\dots\}$.

Let $\phi : \mathcal{A} \rightarrow \mathcal{L}(H)$ be a $*$ -homomorphism. Then $ax := \phi(a)x$ defines a left action of $\mathcal{A}$ on H , and we call H a *C^* -correspondence* over $\mathcal{A}$ (in [23] and [8] a C^* -correspondence is called a Hilbert bimodule, but it now seems that the term Hilbert bimodule has been reserved for a special kind of C^* -correspondences cf. [24]).

A *Toeplitz representation* (ψ, π) of H in a C^* -algebra B consists of a linear map $\psi : H \rightarrow B$ and a $*$ -homomorphism $\pi : \mathcal{A} \rightarrow B$ such that

$$\psi(\xi a) = \psi(\xi)\pi(a), \quad \psi(\xi)^*\psi(\eta) = \pi(\langle \xi, \eta \rangle), \quad \text{and} \quad \psi(a\xi) = \pi(a)\psi(\xi)$$

for $\xi, \eta \in \mathbf{H}$ and $a \in \mathcal{A}$. Given such a representation, there is a homomorphism $\pi^{(1)} : \mathcal{K}(\mathbf{H}) \rightarrow B$ which satisfies

$$\pi^{(1)}(\theta_{\xi, \eta}) = \psi(\xi)\psi(\eta)^*$$

for all $\xi, \eta \in \mathbf{H}$, and we then have

$$\pi^{(1)}(T)\psi(\xi) = \psi(T\xi)$$

for every $T \in \mathcal{K}(\mathbf{H})$ and $\xi \in \mathbf{H}$. If $\rho : B \rightarrow C$ is a $*$ -homomorphism between C^* -algebras, then $(\rho \circ \psi, \rho \circ \pi)$ is a Toeplitz representation of $\mathbf{H}$, and since

$$(\rho \circ \pi)^{(1)}(\theta_{\xi, \eta}) = (\rho \circ \psi(\xi))(\rho \circ \psi(\eta))^* = \rho \circ \pi^{(1)}(\theta_{\xi, \eta})$$

for all $\xi, \eta \in \mathbf{H}$, by linearity and continuity we have

$$(\rho \circ \pi)^{(1)} = \rho \circ \pi^{(1)}.$$

We denote by $\mathcal{J}(\mathbf{H})$ the closed two-sided ideal $\phi^{-1}(\mathcal{K}(\mathbf{H}))$ in $\mathcal{A}$, and we say that a Toeplitz representation (ψ, π) of $\mathbf{H}$ is *Cuntz-Pimsner coinvariant* if

$$\pi^{(1)}(\phi(a)) = \pi(a)$$

for all $a \in \mathcal{J}(\mathbf{H})$.

Theorem 3.1 ([8, Proposition 1.3] (cf. [23] and [24, Proposition 1.6])). *Let $\mathbf{H}$ be a C^* -correspondence over $\mathcal{A}$. Then there is a C^* -algebra $\mathcal{O}_{\mathbf{H}}$ and a Cuntz-Pimsner coinvariant Toeplitz representation $(k_{\mathbf{H}}, k_{\mathcal{A}}) : \mathbf{H} \rightarrow \mathcal{O}_{\mathbf{H}}$ which satisfies:*

1. *For every Cuntz-Pimsner coinvariant Toeplitz representation (ψ, π) of $\mathbf{H}$, there is a homomorphism $\psi \times \pi$ of $\mathcal{O}_{\mathbf{H}}$ such that $(\psi \times \pi) \circ k_{\mathbf{H}} = \psi$ and $(\psi \times \pi) \circ k_{\mathcal{A}} = \pi$,*
2. *$\mathcal{O}_{\mathbf{H}}$ is generated as a C^* -algebra by $k_{\mathbf{H}}(\mathbf{H}) \cup k_{\mathcal{A}}(\mathcal{A})$.*

Remark 3.2. The triple $(\mathcal{O}_{\mathbf{H}}, k_{\mathbf{H}}, k_{\mathcal{A}})$ is unique: if $(\mathcal{X}, k'_{\mathbf{H}}, k'_{\mathcal{A}})$ has similar properties, then there is an isomorphism $\theta : \mathcal{O}_{\mathbf{H}} \rightarrow \mathcal{X}$ such that $\theta \circ k_{\mathbf{H}} = k'_{\mathbf{H}}$ and $\theta \circ k_{\mathcal{A}} = k'_{\mathcal{A}}$. Thus there is a strongly continuous gauge action $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{O}_{\mathbf{H}}$ which satisfies $\gamma_z(k_{\mathcal{A}}(a)) = k_{\mathcal{A}}(a)$ and $\gamma_z(k_{\mathbf{H}}(x)) = zk_{\mathbf{H}}(x)$ for $a \in \mathcal{A}$ and $x \in \mathbf{H}$.

4 C^* -correspondences associated with subshifts

We will now define the C^* -correspondence H_X that we associate to a subshift X .

We start out by defining the C^* -algebra $\tilde{\mathcal{D}}_X$ which H_X is a C^* -correspondence over.

Definition 4.1. For every subshift X we let $\mathfrak{B}(X)$ be the abelian C^* -algebra of all bounded functions on X , and $\tilde{\mathcal{D}}_X$ the C^* -subalgebra of $\mathfrak{B}(X)$ generated by $\{1_{C(\mu, \nu)} \mid \mu, \nu \in \mathfrak{a}^*\}$.

It turns out that the spectrum of $\tilde{\mathcal{D}}_X$ is the right underlying compact space for the C^* -algebra that we are going to associate with subshifts, but since we will not need an explicit description of this compact space we are not going to give one, but instead work with $\tilde{\mathcal{D}}_X$.

Definition 4.2. Let X be a subshift. For every $a \in \mathfrak{a}$ let $\tilde{\mathcal{D}}_a$ be the ideal in $\tilde{\mathcal{D}}_X$ generated by $1_{\sigma(C(a))}$. Let H_X be the right Hilbert $\tilde{\mathcal{D}}_X$ -module

$$\bigoplus_{a \in \mathfrak{a}} \tilde{\mathcal{D}}_a$$

with the right action is given by

$$(f_a)_{a \in \mathfrak{a}} f = (f_a f)_{a \in \mathfrak{a}}$$

and the inner product by

$$\langle (f_a)_{a \in \mathfrak{a}}, (g_a)_{a \in \mathfrak{a}} \rangle = \sum_{a \in \mathfrak{a}} f_a^* g_a$$

for $(f_a)_{a \in \mathfrak{a}}, (g_a)_{a \in \mathfrak{a}} \in \bigoplus_{a \in \mathfrak{a}} \tilde{\mathcal{D}}_a$ and $f \in \tilde{\mathcal{D}}_X$.

Proposition 4.3. Let X be a subshift and let $a \in \mathfrak{a}$. Define a $*$ -homomorphism $\tilde{\lambda}_a : \mathfrak{B}(X) \rightarrow \mathfrak{B}(X)$ by letting

$$\tilde{\lambda}_a(f)(x) = \begin{cases} f(ax) & \text{if } ax \in X \\ 0 & \text{if } ax \notin X \end{cases}$$

for every $f \in \mathfrak{B}(X)$ and every $x \in X$.

Then $\tilde{\lambda}_a(\tilde{\mathcal{D}}_X) \subseteq \tilde{\mathcal{D}}_a$.

Proof. Let $\mu, \nu \in \mathfrak{a}^*$ with $|\nu| \geq 1$. For every $x \in X$ is

$$\begin{aligned}\tilde{\lambda}_a(1_{C(\mu, \nu)})(x) &= \begin{cases} 1_{C(\mu, \nu)}(ax) & \text{if } ax \in X \\ 0 & \text{if } ax \notin X \end{cases} \\ &= \begin{cases} 1 & \text{if } a = \nu_1, x_1 = \nu_2, \dots, x_{|\nu|-1} = \nu_{|\nu|}, \mu\sigma^{|\nu|-1}(x) \in X, ax \in X \\ 0 & \text{else.} \end{cases}\end{aligned}$$

So $\tilde{\lambda}_a(1_{C(\mu, \nu)}) = 0$ if $a \neq \nu_1$, and

$$\tilde{\lambda}_a(1_{C(\mu, \nu)}) = 1_{C(\mu, \nu_2 \nu_3 \dots \nu_{|\nu|})} 1_{\sigma(C(a))}$$

if $a = \nu_1$. Hence $\tilde{\lambda}_a(1_{C(\nu, \mu)}) \in \tilde{D}_a$. In a similar way, we see that $\tilde{\lambda}_a(1_{C(\mu, \epsilon)}) = 1_{C(a\mu, \epsilon)}$, so $\tilde{\lambda}_a(1_{C(\nu, \epsilon)}) \in \tilde{D}_a$. Thus $\tilde{\lambda}_a(\tilde{D}_X) \subseteq \tilde{D}_a$, since $\tilde{D}_X$ is generated by $\{1_{C(\mu, \nu)} \mid \mu, \nu \in \mathfrak{a}^*\}$. $\square$

Definition 4.4. Let X be a subshift. We let $\phi : \tilde{D}_X \rightarrow \mathcal{L}(H_X)$ be the $*$ -homomorphism defined by

$$\phi(f)((f_a)_{a \in \mathfrak{a}}) = (\tilde{\lambda}_a(f)f_a)_{a \in \mathfrak{a}}$$

for every $f \in \tilde{D}_X$ and every $(f_a)_{a \in \mathfrak{a}} \in H_X$. With this H_X becomes a C^* -correspondence.

5 The C^* -algebra associated with a subshift

We are now ready to define the C^* -algebra $\mathcal{O}_X$ associated with a subshift X .

Definition 5.1. Let X be a subshift. The C^* -algebra $\mathcal{O}_X$ associated with X is the C^* -algebra $\mathcal{O}_{H_X}$ from Theorem 3.1, where H_X is the C^* -correspondence defined above.

We will now take a closer look at $\mathcal{O}_X$. First, we show that $\mathcal{O}_X$ is unital.

Lemma 5.2. Let X be a subshift and let $1 = 1_{C(\epsilon, \epsilon)}$ be the unit of $\tilde{D}_X$. Then $k_{\tilde{D}_X}(1)$ is a unit for $\mathcal{O}_X$.

Proof. We have that

$$k_{H_X}(\xi)k_{\tilde{D}_X}(1) = k_{H_X}(\xi 1) = k_{H_X}(\xi),$$

and

$$k_{\tilde{D}_X}(1)k_{H_X}(\xi) = k_{H_X}(\phi(1)\xi) = k_{H_X}(\xi)$$

for every $\xi \in \mathbf{H}_X$. Since we also have that

$$k_{\tilde{\mathcal{D}}_X}(1)k_{\tilde{\mathcal{D}}_X}(f) = k_{\tilde{\mathcal{D}}_X}(f)k_{\tilde{\mathcal{D}}_X}(1) = k_{\tilde{\mathcal{D}}_X}(f)$$

for every $f \in \tilde{\mathcal{D}}_X$, and $\mathcal{O}_X$ is generated by $k_{\mathbf{H}_X}(\mathbf{H}_X) \cup k_{\tilde{\mathcal{D}}_X}(\tilde{\mathcal{D}}_X)$, we have that $k_{\tilde{\mathcal{D}}_X}(1)$ is a unit for $\mathcal{O}_X$. $\square$

We will denote the unit of $\mathcal{O}_X$ by I .

Definition 5.3. Let X be a subshift. For every $a \in \mathfrak{a}$ let ξ_a be the element $(f_{a'})_{a' \in \mathfrak{a}} \in \mathbf{H}_X$ where $f_a = 1_{\sigma(C(a))}$ and $f_{a'} = 0$ for $a' \neq a$, and let for every $\mu \in \mathfrak{a}^*$, S_μ be the product $k_{\mathbf{H}_X}(\xi_{\mu_1})k_{\mathbf{H}_X}(\xi_{\mu_2}) \cdots k_{\mathbf{H}_X}(\xi_{\mu_{|\mu|}}) \in \mathcal{O}_X$ with the convention that $S_\epsilon = I$.

Lemma 5.4. Let X be a subshift. Let 1 be the unit of $\tilde{\mathcal{D}}_X$, and Id the unit of $\mathcal{L}(\mathbf{H}'_X)$. Then

$$k_{\tilde{\mathcal{D}}_X}(1) = k_{\tilde{\mathcal{D}}_X}^{(1)}(\text{Id}) = \sum_{a \in \mathfrak{a}} S_a S_a^*$$

is the unit of $\mathcal{O}_X$.

Proof. It is easy to check that

$$\phi(1) = \text{Id} = \sum_{a \in \mathfrak{a}} \theta_{\xi_a, \xi_a^*},$$

so since $(k_{\mathbf{H}_X}, k_{\tilde{\mathcal{D}}_X})$ is a Cuntz-Pimsner coinvariant representation,

$$k_{\tilde{\mathcal{D}}_X}(1) = k_{\tilde{\mathcal{D}}_X}^{(1)}(\text{Id}) = \sum_{a \in \mathfrak{a}} S_a S_a^*,$$

and we know from Lemma 5.2 that $k_{\tilde{\mathcal{D}}_X}(1)$ is the unit of $\mathcal{O}_X$. $\square$

Lemma 5.5. Let X be a subshift. Then

$$k_{\tilde{\mathcal{D}}_X}(1_{C(\mu, \nu)}) = S_\nu S_\mu^* S_\mu S_\nu^*$$

for every $\mu, \nu \in \mathfrak{a}^*$.

Proof. Since

$$\begin{aligned} S_a^* S_a &= k_{\mathbf{H}_X}(\xi_a)^* k_{\mathbf{H}_X}(\xi_a) \\ &= k_{\tilde{\mathcal{D}}_X}(\langle \xi_a, \xi_a \rangle) \\ &= k_{\tilde{\mathcal{D}}_X}(1_{\sigma(C(a))}), \end{aligned}$$

and

$$\begin{aligned}
S_a^* k_{\tilde{\mathcal{D}}_X} \left(1_{\sigma|\mu'| (C(\mu'))} \right) S_a &= k_{\mathbf{H}_X}(\xi_a)^* k_{\mathbf{H}_X} \left(\phi' \left(1_{\sigma|\mu'| (C(\mu'))} \right) \xi_a \right) \\
&= k_{\mathbf{H}_X}(\xi_a)^* k_{\mathbf{H}_X} \left(\xi_a \tilde{\lambda}_a \left(1_{\sigma|\mu'| (C(\mu'))} \right) \right) \\
&= k_{\mathbf{H}_X}(\xi_a)^* k_{\mathbf{H}_X}(\xi_a) k_{\tilde{\mathcal{D}}_X} \left(\tilde{\lambda}_a \left(1_{\sigma|\mu'| (C(\mu'))} \right) \right) \\
&= k_{\tilde{\mathcal{D}}_X} \left(1_{\sigma(C(a))} \tilde{\lambda}_a \left(1_{\sigma|\mu'| (C(\mu'))} \right) \right) \\
&= k_{\tilde{\mathcal{D}}_X} \left(1_{\sigma|\mu'| a| (C(\mu' a))} \right)
\end{aligned}$$

for every $a \in \mathfrak{a}$ and every $\mu' \in \mathfrak{a}^*$, we have that

$$S_\mu^* S_\mu = k_{\tilde{\mathcal{D}}_X} \left(1_{\sigma|\mu| (C(\mu))} \right)$$

for every $\mu \in \mathfrak{a}^*$.

It is easy to check that for every $f \in \tilde{\mathcal{D}}_X$ is

$$\phi(f) = \sum_{a \in \mathfrak{a}} \theta_{\xi_a \tilde{\lambda}_a(f), \xi_a^*}.$$

Let $\mu, \nu \in \mathfrak{a}^*$ with $|\nu| \geq 1$ and $a \in \mathfrak{a}$. Then as proved in the proof of Proposition 4.3

$$\tilde{\lambda}_a (1_{C(\mu, \nu)}) = 0$$

if $a \neq \nu_1$, and

$$\tilde{\lambda}_a (1_{C(\mu, \nu)}) = 1_{C(\mu, \nu_2, \dots, \nu_{|\nu|})} 1_{\sigma(C(a))}$$

if $a = \nu_1$.

So

$$\begin{aligned}
k_{\tilde{\mathcal{D}}_X} (1_{C(\mu, \nu)}) &= k_{\tilde{\mathcal{D}}_X}^{(1)} (\phi (1_{C(\mu, \nu)})) \\
&= k_{\tilde{\mathcal{D}}_X}^{(1)} \left(\theta_{\xi_{\nu_1} 1_{C(\mu, \nu_2, \dots, \nu_{|\nu|})}, \xi_{\nu_1}} \right) \\
&= k_{\mathbf{H}_X}(\xi_{\nu_1} 1_{C(\mu, \nu_2, \dots, \nu_{|\nu|})}) k_{\mathbf{H}_X}(\xi_{\nu_1})^* \\
&= S_{\nu_1} k_{\tilde{\mathcal{D}}_X} (1_{C(\mu, \nu_2, \dots, \nu_{|\nu|})}) S_{\nu_1}^*.
\end{aligned}$$

Hence

$$k_{\tilde{\mathcal{D}}_X} (1_{C(\mu, \nu)}) = S_\nu S_\mu^* S_\mu S_\nu^*$$

for all $\mu, \nu \in \mathfrak{a}^*$. □

Proposition 5.6. *Let X be a subshift. Then $\mathcal{O}_X$ is generated by $\{S_a\}_{a \in \mathfrak{a}}$.*

Proof. $\mathcal{O}_X$ is by Theorem 3.1 generated by $k_{H_X}(H_X) \cup k_{\tilde{\mathcal{D}}_X}(\tilde{\mathcal{D}}_X)$.

First notice that $k_{\tilde{\mathcal{D}}_X}(1) = \sum_{a \in \mathfrak{a}} S_a S_a^*$ is in the C^* -algebra generated by $\{S_a\}_{a \in \mathfrak{a}}$. Since

$$k_{\tilde{\mathcal{D}}_X}(1_{C(\mu, \nu)}) = S_\nu S_\mu^* S_\mu S_\nu^*$$

for all $\mu, \nu \in \mathfrak{a}^*$, and $\tilde{\mathcal{D}}_X$ is generated by $\{1_{C(\mu, \nu)} \mid \mu, \nu \in \mathfrak{a}^*\}$, we have that $k_{\tilde{\mathcal{D}}_X}(\tilde{\mathcal{D}}_X)$ is in the C^* -algebra generated by $\{S_a\}_{a \in \mathfrak{a}}$.

Let $(f_a)_{a \in \mathfrak{a}} \in H_X$. Then

$$(f_a)_{a \in \mathfrak{a}} = \sum_{a \in \mathfrak{a}} \xi_a f_a,$$

so $k_{H_X}((f_a)_{a \in \mathfrak{a}}) = \sum_{a \in \mathfrak{a}} S_a k_{\tilde{\mathcal{D}}_X}(f_a)$, and $k_{H_X}((f_a)_{a \in \mathfrak{a}})$ is in the C^* -algebra generated by $\{S_a\}_{a \in \mathfrak{a}}$. Hence $\mathcal{O}_X$ is generated by $\{S_a\}_{a \in \mathfrak{a}}$. $\square$

6 The structure of C^* -algebras generated by partial isometries

We have now established that $\mathcal{O}_X$ is a unital C^* -algebra generated by partial isometries $\{S_a\}_{a \in \mathfrak{a}}$, which by Lemma 5.4 and 5.5 satisfy

$$\sum_{a \in \mathfrak{a}} S_a S_a^* = I, \tag{1}$$

$$S_\mu^* S_\mu S_\nu S_\nu^* = S_\nu S_\nu^* S_\mu^* S_\mu, \tag{2}$$

$$S_\mu^* S_\mu S_\nu^* S_\nu = S_\nu^* S_\nu S_\mu^* S_\mu, \tag{3}$$

where $S_\mu = S_{\mu_1} \cdots S_{\mu_{|\mu|}}$ and $S_\nu = S_{\nu_1} \cdots S_{\nu_{|\nu|}}$, for every $\mu, \nu \in \mathfrak{a}^*$.

We will now take a closer look at unital C^* -algebras generated by partial isometries $\{S_a\}_{a \in \mathfrak{a}}$ that satisfy the 3 relations above.

So in the rest of this section, $\mathfrak{a}$ will be an alphabet and $\mathcal{O}$ will be a unital C^* -algebra generated by partial isometries $\{S_a\}_{a \in \mathfrak{a}}$, which satisfy the relations (1), (2) and (3) above.

Lemma 6.1. *For every $\mu \in \mathfrak{a}^*$, S_μ is a partial isometry.*

Proof. We will prove the lemma by induction over the length of $|\mu|$. If $|\mu| = 1$, then S_μ is a partial isometry by definition. Assume now that S_ν is

a partial isometry and $a \in \mathfrak{a}$. Then

$$\begin{aligned}
S_{\nu a} S_{\nu a}^* S_{\nu a} &= S_{\nu} S_a S_a^* S_{\nu}^* S_{\nu} S_a \\
&= S_{\nu} S_{\nu}^* S_{\nu} S_a S_a^* S_a \\
&= S_{\nu} S_a \\
&= S_{\nu a}.
\end{aligned}$$

So $S_{\nu a}$ is a partial isometry. Hence S_{μ} is a partial isometry for every $\mu \in \mathfrak{a}^*$. $\square$

For $\mu \in \mathfrak{a}^*$ we set $A_{\mu} = S_{\mu}^* S_{\mu}$. We notice that since $\sum_{a \in \mathfrak{a}} S_a S_a^* = I$, the projections $\{S_a S_a^*\}_{a \in \mathfrak{a}}$ are mutually orthogonal, so $S_a S_a^* S_b S_b^* = 0$ for $a \neq b$.

Lemma 6.2. *Let $\mu, \nu \in \mathfrak{a}^*$ with $|\mu| = |\nu|$. If $S_{\mu}^* S_{\nu} \neq 0$, then $\mu = \nu$ and $S_{\mu}^* S_{\nu} = A_{\mu}$.*

Proof. We will prove the lemma by induction over the length of μ and ν . If the length is 1 and $\mu \neq \nu$, then

$$\begin{aligned}
S_{\mu}^* S_{\nu} &= S_{\mu}^* S_{\mu} S_{\mu}^* S_{\nu} S_{\nu}^* S_{\nu} \\
&= 0
\end{aligned}$$

since $S_{\mu} S_{\mu}^* S_{\nu} S_{\nu}^* = 0$. So since $S_{\mu}^* S_{\nu} \neq 0$, we have that $\mu = \nu$ and $S_{\mu}^* S_{\nu} = A_{\mu}$.

Now assume that we have proved the lemma in case $|\mu| = |\nu| = n$, and assume that $|\mu'| = |\nu'| = n+1$ and $S_{\mu'}^* S_{\nu'} \neq 0$. Set $\mu = (\mu'_1, \dots, \mu'_n)$ and $\nu = (\nu'_1, \dots, \nu'_n)$. Then $S_{\mu}^* S_{\nu} \neq 0$, so $\mu = \nu$. Since

$$\begin{aligned}
0 &\neq S_{\mu'}^* S_{\nu'} \\
&= S_{\mu'_{n+1}}^* S_{\mu}^* S_{\nu} S_{\nu'_{n+1}} \\
&= S_{\mu'_{n+1}}^* S_{\mu'_{n+1}} S_{\mu'_{n+1}}^* S_{\mu}^* S_{\mu} S_{\nu'_{n+1}} S_{\nu'_{n+1}}^* S_{\nu'_{n+1}} \\
&= S_{\mu'_{n+1}}^* S_{\mu'_{n+1}} S_{\mu'_{n+1}}^* S_{\nu'_{n+1}} S_{\nu'_{n+1}}^* S_{\mu}^* S_{\mu} S_{\nu'_{n+1}},
\end{aligned}$$

we have that $\mu'_{n+1} = \nu'_{n+1}$, and hence $\mu' = \nu'$. So the lemma is true. $\square$

For each $l \in \mathbb{N}_0$ we denote by $\mathcal{A}_l(\mathcal{O})$ the C^* -subalgebra of $\mathcal{O}$ generated by $\{A_{\mu}\}_{\mu \in \mathfrak{a}_l}$. Since $\mathcal{A}_l(\mathcal{O})$ is generated by a finite number of mutually commuting projection, there exist a finite number of mutually orthogonal projections E_i^l , $i = 1, \dots, m(l)$, such that $(E_i^l)_{i=1, \dots, m(l)}$ is a basis for $\mathcal{A}_l(\mathcal{O})$. We have that $\overline{\bigcup_{l \in \mathbb{N}_0} \mathcal{A}_l(\mathcal{O})}$ is the C^* -algebra generated by $\{A_{\mu}\}_{\mu \in \mathfrak{a}^*}$. We denoted this C^* -algebra by $\mathcal{A}(\mathcal{O})$. Since $\mathcal{A}_l(\mathcal{O})$ is finite dimensional and $\mathcal{A}_l(\mathcal{O}) \subseteq \mathcal{A}_{l+1}(\mathcal{O})$ for every $l \in \mathbb{N}_0$, $\mathcal{A}(\mathcal{O})$ is an AF-algebra.

Lemma 6.3. For $1 \leq k \leq l$, $\mu \in \mathfrak{a}^k$ and $i \in \{1, 2, \dots, m(l)\}$, the following two conditions are equivalent:

a) $S_\mu E_i^l S_\mu^* \neq 0$,

b) $A_\mu E_i^l \neq 0$.

Proof. Since

$$S_\mu E_i^l S_\mu^* = S_\mu A_\mu E_i^l S_\mu^*,$$

and

$$A_\mu E_i^l = S_\mu^* S_\mu E_i^l S_\mu^* S_\mu,$$

we have that

$$S_\mu E_i^l S_\mu^* \neq 0 \Leftrightarrow A_\mu E_i^l \neq 0.$$

□

Lemma 6.4. Let $1 \leq k \leq l$. Then

a) For $i, i' \in \{1, 2, \dots, m(l)\}$ and $\mu, \mu' \in \mathfrak{a}^k$ is

$$S_\mu E_i^l S_\mu^* S_{\mu'} E_{i'}^l S_{\mu'}^* = \begin{cases} S_\mu E_i^l S_\mu^* & \text{if } \mu = \mu' \text{ and } i = i' \\ 0 & \text{if } \mu \neq \mu' \text{ or } i \neq i'. \end{cases}$$

b) $(S_\mu E_i^l S_\mu^*)^* = S_\mu E_i^l S_\mu^*$ for $i \in \{1, 2, \dots, m(l)\}$ and $\mu \in \mathfrak{a}^k$.

Proof. a): By Lemma 6.2

$$\begin{aligned} S_\mu E_i^l S_\mu^* S_{\mu'} E_{i'}^l S_{\mu'}^* &= \begin{cases} S_\mu E_i^l A_\mu E_{i'}^l S_{\mu'}^* & \text{if } \mu = \mu' \\ 0 & \text{if } \mu \neq \mu' \end{cases} \\ &= \begin{cases} S_\mu A_\mu E_i^l E_{i'}^l S_{\mu'}^* & \text{if } \mu = \mu' \\ 0 & \text{if } \mu \neq \mu' \end{cases} \\ &= \begin{cases} S_\mu E_i^l S_{\mu'}^* & \text{if } \mu = \mu' \text{ and } i = i' \\ 0 & \text{if } \mu \neq \mu' \text{ or } i \neq i'. \end{cases} \end{aligned}$$

b): Obviously.

□

7 The universal property of $\mathcal{O}_X$

We let $\tilde{\mathcal{A}}_X$ be the C^* -subalgebra of $\tilde{\mathcal{D}}_X$ generated by $\{1_{\sigma|\mu|(C(\mu))} \mid \mu \in \mathfrak{a}^*\}$.

Lemma 7.1. *Let X be a subshift, $\mathcal{X}$ a C^* -algebra, $\psi : \tilde{\mathcal{A}}_X \rightarrow \mathcal{X}$ a $*$ -homomorphism and $\{S_a\}_{a \in \mathfrak{a}}$ partial isometries in $\mathcal{X}$ such that*

- a) $\sum_{a \in \mathfrak{a}} S_a S_a^* = \psi(1)$,
- b) $S_\mu^* S_\mu S_\nu S_\nu^* = S_\nu S_\nu^* S_\mu^* S_\mu$,
- c) $S_\mu^* S_\mu = \psi\left(1_{\sigma|\mu|(C(\mu))}\right)$,

where $S_\mu = S_{\mu_1} S_{\mu_2} \cdots S_{\mu_{|\mu|}}$ and $S_\nu = S_{\nu_1} S_{\nu_2} \cdots S_{\nu_{|\nu|}}$, for every $\mu, \nu \in \mathfrak{a}^*$.

Then ψ extends to a $*$ -homomorphism from $\tilde{\mathcal{D}}_X$ to $\mathcal{X}$, such that

$$\psi\left(1_{C(\mu, \nu)}\right) = S_\nu S_\nu^* S_\mu^* S_\mu$$

for every $\mu, \nu \in \mathfrak{a}^*$.

Proof. Let $\mathcal{O}$ be the C^* -subalgebra of $\mathcal{X}$ generated by $\{S_a\}_{a \in \mathfrak{a}}$. Since $S_\mu^* S_\mu = \psi\left(1_{\sigma|\mu|(C(\mu))}\right)$ and $S_\nu^* S_\nu = \psi\left(1_{\sigma|\nu|(C(\nu))}\right)$, we have that $S_\mu^* S_\mu S_\nu^* S_\nu = S_\nu^* S_\nu S_\mu^* S_\mu$ for every $\mu, \nu \in \mathfrak{a}^*$. Since

$$\begin{aligned} S_a \psi(1) &= S_a S_a^* S_a \psi(1) \\ &= S_a \psi\left(1_{C(a)}\right) \psi(1) \\ &= S_a \psi\left(1_{C(a)}\right) \\ &= S_a S_a^* S_a \\ &= S_a \end{aligned}$$

and

$$\begin{aligned} \psi(1) S_a &= \psi(1) S_a S_a^* S_a \\ &= \sum_{a' \in \mathfrak{a}} S_{a'} S_{a'}^* S_a S_a^* S_a \\ &= S_a S_a^* S_a \\ &= S_a \end{aligned}$$

for every $a \in \mathfrak{a}$, $\psi(1)$ is a unit for $\mathcal{O}$. Hence $\mathcal{O}$ is generated by partial isometries $\{S_a\}_{a \in \mathfrak{a}}$ which satisfy the relations (1), (2) and (3) of section 6.

For each $l \in \mathbb{N}_0$, denote by $\tilde{\mathcal{A}}_l$ the C^* -subalgebra of $\tilde{\mathcal{A}}_{\mathbf{X}}$ generated by $\{1_{\sigma^{|\mu|}(C(\mu))} \mid \mu \in \mathfrak{a}_l\}$. Since $\tilde{\mathcal{A}}_l$ is generated by a finite number of mutually commuting projections, there exists a finite number $m(l)$ of mutually disjoint subsets $\mathcal{E}_i^l$, $i = 1, 2, \dots, m(l)$ of $\mathbf{X}$ such that

$$\left\{1_{\mathcal{E}_i^l} \mid i \in \{1, 2, \dots, m(l)\}\right\}$$

is a basis for $\tilde{\mathcal{A}}_l$.

Then $\psi(1_{\mathcal{E}_i^l})$, $i = 1, 2, \dots, m(l)$ are mutually orthogonal projections in $\mathcal{O}$ and $\text{span}\left\{\psi(1_{\mathcal{E}_i^l}) \mid i \in \{1, 2, \dots, m(l)\}\right\} = \mathcal{A}_l(\mathcal{O})$. So by Lemma 6.4 we have that for $1 \leq k \leq l$ are $S_\nu \psi(1_{\mathcal{E}_i^l}) S_\nu^*$, $i = 1, 2, \dots, m(l)$ mutually orthogonal projections in $\mathcal{O}$.

For each $1 \leq k \leq l$ denote by $\tilde{\mathcal{D}}_k^l$ the C^* -subalgebra of $\tilde{\mathcal{D}}_{\mathbf{X}}$ generated by $\{1_{C(\mu, \nu)} \mid \nu \in \mathfrak{a}^k, \mu \in \mathfrak{a}_l\}$. It is easy to check that

$$1_{C(\nu) \cap \sigma^{-|\nu|}(\mathcal{E}_i^l)}, \quad \nu \in \mathfrak{a}^k, \quad i = 1, 2, \dots, m(l)$$

are mutually orthogonal projections in $\tilde{\mathcal{D}}_k^l$, and since

$$\begin{aligned} 1_{C(\nu) \cap \sigma^{-|\nu|}(\mathcal{E}_i^l)} = 0 &\Rightarrow 1_{\sigma^{|\nu|}(C(\nu))} 1_{\mathcal{E}_i^l} = 0 \\ &\Rightarrow S_\nu^* S_\nu \psi(1_{\mathcal{E}_i^l}) = 0 \\ &\Rightarrow S_\nu \psi(1_{\mathcal{E}_i^l}) S_\nu^* = 0, \end{aligned}$$

there exists a $*$ -homomorphism $\psi_k^l : \tilde{\mathcal{D}}_k^l \rightarrow \mathcal{X}$ such that $\psi_k^l(1_{C(\nu) \cap \sigma^{-|\nu|}(\mathcal{E}_i^l)}) = S_\nu \psi(1_{\mathcal{E}_i^l}) S_\nu^*$ for every $\nu \in \mathfrak{a}^k$ and every $i \in \{1, 2, \dots, m(l)\}$ and hence $\psi_k^l(1_{C(\mu, \nu)}) = S_\nu S_\mu^* S_\mu S_\nu^*$ for every $\nu \in \mathfrak{a}^k$ and every $\mu \in \mathfrak{a}_l$.

For every $k \in \mathbb{N}_0$ denote by $\tilde{\mathcal{D}}_k$ the C^* -subalgebra of $\tilde{\mathcal{D}}_{\mathbf{X}}$ generated by $\{1_{C(\mu, \nu)} \mid \nu \in \mathfrak{a}^k, \mu \in \mathfrak{a}^*\}$. Then

$$\tilde{\mathcal{D}}_k = \overline{\bigcup_{l \geq k} \tilde{\mathcal{D}}_k^l}.$$

Let ι_k^l denote the inclusion of $\tilde{\mathcal{D}}_k^l$ into $\tilde{\mathcal{D}}_k^{l+1}$. Since $\psi_k^{l+1} \circ \iota_k^l = \psi_k^l$ for every $l \geq k$, the ψ_k^l 's induce a $*$ -homomorphism $\psi_k : \tilde{\mathcal{D}}_k \rightarrow \mathcal{O}$ such that $\psi_k(1_{C(\mu, \nu)}) = S_\nu S_\mu^* S_\mu S_\nu^*$ for every $\nu \in \mathfrak{a}^k$ and every $\mu \in \mathfrak{a}^*$.

Since

$$C(\mu, \nu) = \bigcup_{a \in \mathfrak{a}} C(\mu a, \nu a)$$

for every $\mu, \nu \in \mathfrak{a}^*$, $\tilde{\mathcal{D}}_k \subseteq \tilde{\mathcal{D}}_{k+1}$ for every $k \in \mathbb{N}_0$ and the inclusion ι_k of $\tilde{\mathcal{D}}_k$ into $\tilde{\mathcal{D}}_{k+1}$ is given by

$$\iota_k(1_{C(\mu, \nu)}) = \sum_{a \in \mathfrak{a}} 1_{C(\mu a, \nu a)}.$$

Hence $\psi_{k+1} \circ \iota_k = \psi_k$ and since $\tilde{\mathcal{D}}_{\mathbf{X}} = \overline{\bigcup_{k \in \mathbb{N}_0} \tilde{\mathcal{D}}_k}$, the ψ_k 's induce a $*$ -homomorphism $\psi : \tilde{\mathcal{D}}_{\mathbf{X}} \rightarrow \mathcal{O} \subseteq \mathcal{X}$ such that

$$\psi(1_{C(\mu, \nu)}) = S_\nu S_\mu^* S_\mu S_\nu^*$$

for every $\mu, \nu \in \mathfrak{a}^*$. □

We are now ready to state and prove the universal property of $\mathcal{O}_{\mathbf{X}}$.

Theorem 7.2. *Let $\mathbf{X}$ be a subshift. Then $\mathcal{O}_{\mathbf{X}}$ is the universal unital C^* -algebra generated by partial isometries $\{S_a\}_{a \in \mathfrak{a}}$ satisfying*

- a) $\sum_{a \in \mathfrak{a}} S_a S_a^* = I$,
- b) $S_\mu^* S_\mu S_\nu S_\nu^* = S_\nu S_\nu^* S_\mu^* S_\mu$,
- c) *the map $1_{C(\mu)} \mapsto S_\mu^* S_\mu$ extends to a unital $*$ -homomorphism from $\tilde{\mathcal{A}}_{\mathbf{X}}$ to the C^* -algebra generated by $\{S_a\}_{a \in \mathfrak{a}}$,*

where $S_\mu = S_{\mu_1} \cdots S_{\mu_{|\mu|}}$ and $S_\nu = S_{\nu_1} \cdots S_{\nu_{|\nu|}}$ for every $\mu, \nu \in \mathfrak{a}^*$.

Proof. It follows from Lemma 5.4 and 5.5 and Proposition 5.6 together with the fact that $\tilde{\mathcal{A}}_{\mathbf{X}}$ is a C^* -subalgebra of $\tilde{\mathcal{D}}_{\mathbf{X}}$, that $\mathcal{O}_{\mathbf{X}}$ is generated by partial isometries $\{S_a\}_{a \in \mathfrak{a}}$ satisfying a), b) and c).

Assume now that $\mathcal{X}$ is a unital C^* -algebra generated by partial isometries $\{T_a\}_{a \in \mathfrak{a}}$ and that $\pi : \tilde{\mathcal{A}}_{\mathbf{X}} \rightarrow \mathcal{X}$ is a unital $*$ -homomorphism such that

- a) $\sum_{a \in \mathfrak{a}} T_a T_a^* = I$,
- b) $T_\mu^* T_\mu T_\nu T_\nu^* = T_\nu T_\nu^* T_\mu^* T_\mu$,
- c) $T_\mu^* T_\mu = \pi \left(1_{\sigma^{|\mu|}(C(\mu))} \right)$,

where $T_\mu = T_{\mu_1} T_{\mu_2} \cdots T_{\mu_{|\mu|}}$ and $T_\nu = T_{\nu_1} T_{\nu_2} \cdots T_{\nu_{|\nu|}}$, for every $\mu, \nu \in \mathfrak{a}^*$.

By Lemma 7.1, π extends to a $*$ -homomorphism from $\tilde{\mathcal{D}}_{\mathbf{X}}$ to $\mathcal{X}$, such that

$$\pi(1_{C(\mu, \nu)}) = T_\nu T_\mu^* T_\mu T_\nu^*$$

for every $\mu, \nu \in \mathfrak{a}^*$. Let

$$\psi((f_a)_{a \in \mathfrak{a}}) = \sum_{a \in \mathfrak{a}} T_a \pi(f_a)$$

for every $(f_a)_{a \in \mathfrak{a}} \in H_X$. We will show that (ψ, π) is a Cuntz-Pimsner coinvariant representation of H_X .

It is easy to check that $\alpha\psi(\xi) + \beta\psi(\zeta) = \psi(\alpha\xi + \beta\zeta)$ for every $\alpha, \beta \in \mathbb{C}$ and every $\xi, \zeta \in H_X$, and $\psi(\xi)\pi(f) = \psi(\xi f)$ for every $\xi \in H_X$ and every $f \in \tilde{\mathcal{A}}_X$.

Recall from the proof of Proposition 4.3 that for $\mu, \nu \in \mathfrak{a}^*$ with $|\nu| \geq 1$ is $\tilde{\lambda}_a(1_{C(\mu, \nu)}) = 0$ if $a \neq \nu_1$, and $\tilde{\lambda}_a(1_{C(\mu, \nu)}) = 1_{C(\mu, \nu_2 \nu_3 \dots \nu_{|\nu|})} 1_{\sigma(C(a))}$ if $a = \nu_1$. Thus

$$\begin{aligned} \pi(1_{C(\mu, \nu)}) \psi((f_a)_{a \in \mathfrak{a}}) &= \pi(1_{C(\mu, \nu)}) \sum_{a \in \mathfrak{a}} T_a \pi(f_a) \\ &= \sum_{a \in \mathfrak{a}} T_\nu T_\mu^* T_\mu T_\nu^* T_a \pi(f_a) \\ &= T_\nu T_\mu^* T_\mu T_\nu^* T_{\nu_1} \pi(f_{\nu_1}) \\ &= T_{\nu_1} T_{\nu_2 \nu_3 \dots \nu_{|\nu|}} T_\mu^* T_\mu T_{\nu_2 \nu_3 \dots \nu_{|\nu|}}^* \pi(f_{\nu_1}) \\ &= T_{\nu_1} \pi(1_{C(\mu, \nu_2 \nu_3 \dots \nu_{|\nu|})}) \pi(f_{\nu_1}) \\ &= \psi(\xi_{\nu_1} 1_{C(\mu, \nu_2 \nu_3 \dots \nu_{|\nu|})} f_{\nu_1}) \\ &= \psi(\tilde{\lambda}_a(1_{C(\mu, \nu)}) f_a)_{a \in \mathfrak{a}} \\ &= \psi(\phi(1_{C(\mu, \nu)})(f_a)_{a \in \mathfrak{a}}), \end{aligned}$$

for $(f_a)_{a \in \mathfrak{a}} \in H_X$. We also have that

$$\begin{aligned}
\pi \left(1_{\sigma^{|\mu|}(C(\mu))} \right) \psi((f_a)_{a \in \mathfrak{a}}) &= \pi \left(1_{\sigma^{|\mu|}(C(\mu))} \right) \sum_{a \in \mathfrak{a}} T_a \pi(f_a) \\
&= \sum_{a \in \mathfrak{a}} T_\mu^* T_\mu T_a \pi(f_a) \\
&= \sum_{a \in \mathfrak{a}} T_a T_{\mu a}^* T_{\mu a} \pi(f_a) \\
&= \sum_{a \in \mathfrak{a}} T_a \pi \left(1_{\sigma^{|\mu a|}(C(\mu a))} \right) \pi(f_a) \\
&= \psi \left((1_{\sigma^{|\mu a|}(C(\mu a))} f_a)_{a \in \mathfrak{a}} \right) \\
&= \psi \left((\tilde{\lambda}_a (1_{\sigma^{|\mu|}(C(\mu))}) f_a)_{a \in \mathfrak{a}} \right) \\
&= \psi \left(\phi(1_{\sigma^{|\mu|}(C(\mu))}) (f_a)_{a \in \mathfrak{a}} \right).
\end{aligned}$$

Since $\tilde{D}_X$ is generated by $1_{C(\mu, \nu)}$, $\mu, \nu \in \mathfrak{a}^*$, we have that

$$\pi(f) \psi((f_a)_{a \in \mathfrak{a}}) = \psi(\phi(f)(f_a)_{a \in \mathfrak{a}})$$

for every $f \in \tilde{D}_X$ and every $(f_a)_{a \in \mathfrak{a}} \in H_X$.

Since $\sum_{a \in \mathfrak{a}} T_a T_a^* = I$, the projections $\{T_a T_a^*\}_{a \in \mathfrak{a}}$ are mutually orthogonal, so

$$\begin{aligned}
T_a^* T_{a'} &= T_a^* T_a T_a^* T_{a'} T_{a'}^* T_{a'} \\
&= 0
\end{aligned}$$

if $a \neq a'$. Thus

$$\begin{aligned}
\psi((f_a)_{a \in \mathfrak{a}})^* \psi((g_a)_{a \in \mathfrak{a}}) &= \sum_{a \in \mathfrak{a}} \pi(f_a^*) T_a^* \sum_{a' \in \mathfrak{a}} T_{a'} \pi(g_{a'}) \\
&= \sum_{a \in \mathfrak{a}} \pi(f_a^*) T_a^* T_a \pi(g_a) \\
&= \sum_{a \in \mathfrak{a}} \pi(f_a^*) \pi(1_{\sigma(C(a))}) \pi(g_a) \\
&= \pi(\langle (f_a)_{a \in \mathfrak{a}}, (g_a)_{a \in \mathfrak{a}} \rangle)
\end{aligned}$$

for every $(f_a)_{a \in \mathfrak{a}}, (g_a)_{a \in \mathfrak{a}} \in H_X$.

Finally we see that for every $f \in \tilde{\mathcal{D}}_{\mathbf{X}}$ is $\phi(f) = \sum_{a \in \mathfrak{a}} \theta_{\xi_a \tilde{\lambda}_a(f), \xi_a}$, so

$$\begin{aligned} \pi^{(1)}(\phi(f)) &= \sum_{a \in \mathfrak{a}} \psi(\xi_a \tilde{\lambda}_a(f)) \psi(\xi_a)^* \\ &= \sum_{a \in \mathfrak{a}} T_a \pi(\tilde{\lambda}_a(f)) T_a^* \\ &= \sum_{a \in \mathfrak{a}} \pi(f) T_a T_a^* \\ &= \pi(f). \end{aligned}$$

Thus (ψ, π) is a Cuntz-Pimsner coinvariant representation of $\mathbf{H}_{\mathbf{X}}$, so it follows from Theorem 3.1 that there exists a $*$ -homomorphism $\psi \times \pi$ from $\mathcal{O}_{\mathbf{X}}$ to $\mathcal{X}$ such that $\psi \times \pi(k_{\mathbf{H}_{\mathbf{X}}}((f_a)_{a \in \mathfrak{a}})) = \psi((f_a)_{a \in \mathfrak{a}})$ for every $(f_a)_{a \in \mathfrak{a}} \in \mathbf{H}_{\mathbf{X}}$ and hence

$$\psi \times \pi(S_a) = \psi \times \pi(k_{\mathbf{H}_{\mathbf{X}}}(\xi_a)) = \psi(\xi_a) = T_a$$

for every $a \in \mathfrak{a}$. $\square$

Remark 7.3. It follows from Lemma 7.1 that $\mathcal{O}_{\mathbf{X}}$ also can be characterized as the universal C^* -algebra generated by partial isometries $\{S_a\}_{a \in \mathfrak{a}}$ such that the map $1_{C(\mu, \nu)} \mapsto S_{\nu} S_{\mu}^* S_{\mu} S_{\nu}^*$ extends to a $*$ -homomorphism from $\tilde{\mathcal{D}}_{\mathbf{X}}$ to the C^* -algebra generated by $\{S_a\}_{a \in \mathfrak{a}}$, where $S_{\mu} = S_{\mu_1} \cdots S_{\mu_{|\mu|}}$ and $S_{\nu} = S_{\nu_1} \cdots S_{\nu_{|\nu|}}$ for every $\mu, \nu \in \mathfrak{a}^*$.

Remark 7.4. Condition b) can be replaced by

$$\text{b')} \quad S_{\mu}^* S_{\mu} S_{\nu} = S_{\nu} S_{\mu\nu}^* S_{\mu\nu},$$

because b') implies that $S_{\mu}^* S_{\mu} S_{\nu} S_{\nu}^* = S_{\nu} S_{\mu\nu}^* S_{\mu\nu} S_{\nu}^* = S_{\nu} S_{\nu}^* S_{\mu}^* S_{\mu} S_{\nu} S_{\nu}^*$ and $S_{\nu} S_{\nu}^* S_{\mu}^* S_{\mu} = S_{\nu} (S_{\mu}^* S_{\mu} S_{\nu})^* = S_{\nu} (S_{\nu} S_{\mu\nu}^* S_{\nu\mu})^* = S_{\nu} S_{\nu}^* S_{\mu}^* S_{\mu} S_{\nu} S_{\nu}^*$, and thus $S_{\mu}^* S_{\mu} S_{\nu} S_{\nu}^* = S_{\nu} S_{\nu}^* S_{\mu}^* S_{\mu}$, and b) implies that $S_{\mu}^* S_{\mu} S_{\nu} = S_{\mu}^* S_{\mu} S_{\nu} S_{\nu}^* S_{\nu} = S_{\nu} S_{\nu}^* S_{\mu}^* S_{\mu} S_{\nu} = S_{\nu} S_{\mu\nu}^* S_{\mu\nu}$.

Thus if we for a two-sided subshift Λ define $\mathcal{O}_{\Lambda}$ to be $\mathcal{O}_{\mathbf{X}_{\Lambda}}$ where $\mathbf{X}_{\Lambda}$ is the one-sided subshift of Λ (cf. Section 2), then $\mathcal{O}_{\Lambda}$ has the universal property [13, Theorem 4.9] and also has the right underlying compact space (cf. [17, Lemma 3.1]) and thus satisfy all of the results of [2, 10, 13–21]

Remark 7.5. It is easy to check that for a two-sided subshift Λ , the partial isometries of [5, Definition 2.1] satisfy a), b) and c) of Theorem 7.2 (with $\mathbf{X} = \mathbf{X}_{\Lambda}$, the one-sided subshift of Λ (cf. Section 2)). Thus there is a surjective $*$ -homomorphism from $\mathcal{O}_{\mathbf{X}_{\Lambda}}$ to $\mathcal{O}_{\Lambda}$ of [5]. If Λ satisfies the condition (I) of [5], then this $*$ -homomorphism is an isomorphism. There are examples of subshifts for which the $*$ -homomorphism is not injective (an example of this is if Λ only consists of one point).

Remark 7.6. In [6] a C^* -algebra $\mathcal{D}_X \rtimes_{\alpha, \mathcal{L}} \mathbb{N}$ has for every one-sided subshift X been constructed by using Exel's crossed product of a C^* -algebra of an endomorphism. It follows from [6, Remark 9] that $\mathcal{O}_X$ and $\mathcal{D}_X \rtimes_{\alpha, \mathcal{L}} \mathbb{N}$ are isomorphic for every one-sided subshift X .

Remark 7.7. It follows from Theorem 10 of Chapter 2 of [3] that $\mathcal{O}_X$ can be constructed as the C^* -algebra of a groupoid.

8 $\mathcal{O}_X$ is an invariant

We will now show that $\mathcal{O}_X$ is an invariant for subshifts. We will do that by showing that if two subshifts X and Y are conjugate, then $\mathcal{H}_X$ and $\mathcal{H}_Y$ are isomorphic as C^* -correspondences, and it then follows that $\mathcal{O}_X$ and $\mathcal{O}_Y$ are isomorphic.

Definition 8.1. Let $\mathcal{X}$ and $\mathcal{X}'$ be C^* -algebras, (H, ϕ) a C^* -correspondence over $\mathcal{X}$ and (H, ϕ') a C^* -correspondence over $\mathcal{X}'$. If there exist a $*$ -isomorphism $\psi : \mathcal{X} \rightarrow \mathcal{X}'$ and a bijective map $T : H \rightarrow H$ such that

$$\langle T\xi, \zeta \rangle = \psi(\langle \xi, T^{-1}\zeta \rangle),$$

and

$$T(\phi(X)\xi) = \phi'(\psi(X))(T\xi)$$

for all $\xi \in H$, $\zeta \in H'$, $X \in \mathcal{X}$; then we say that (T, ψ) is an C^* -correspondence isomorphism, (H, ϕ) and (H', ϕ') are isomorphic and we write $H \cong H'$.

It easily follows from Theorem 3.1 that if (T, ψ) is an C^* -correspondence isomorphism from (H, ϕ) to (H', ϕ') , then there exists a $*$ -isomorphism $T \times \psi$ from $\mathcal{O}_H$ to $\mathcal{O}_{H'}$ such that $T \times \psi \circ k_H = k_{H'} \circ T$ and $T \times \psi \circ k_{\mathcal{X}} = k_{\mathcal{X}'} \circ \psi$.

Lemma 8.2. Let X be a subshift. Define a $*$ -homomorphism $\tilde{\phi}_X : \mathfrak{B}(X) \rightarrow \mathfrak{B}(X)$ by letting

$$\tilde{\phi}_X(f)(x) = f(\sigma(x))$$

for every $f \in \mathfrak{B}(X)$ and every $x \in X$.

Then $\tilde{\phi}_X(\tilde{\mathcal{D}}_X) \subseteq \tilde{\mathcal{D}}_X$.

Proof. Let $\mu, \nu \in \mathfrak{a}^*$. Then

$$\sigma^{-1}(C(\mu, \nu)) = \bigcup_{a \in \mathfrak{a}} C(\mu, a\nu),$$

so

$$\begin{aligned}
\tilde{\phi}_X(1_{C(\mu,\nu)}) &= 1_{\sigma^{-1}(C(\mu,\nu))} \\
&= 1_{\bigcup_{a \in \mathfrak{a}} C(\mu,a\nu)} \\
&= \sum_{a \in \mathfrak{a}} 1_{C(\mu,a\nu)} \in \tilde{\mathcal{D}}_X.
\end{aligned}$$

Thus, since $\tilde{\mathcal{D}}_X$ is generated by $\{1_{C(\mu,\nu)} \mid \mu, \nu \in \mathfrak{a}^*\}$ and $\tilde{\phi}_X$ is a $*$ -homomorphism, it follows that $\tilde{\phi}_X(\tilde{\mathcal{D}}_X) \subseteq \tilde{\mathcal{D}}_X$. $\square$

Lemma 8.3. *Let X be a subshift. Then we have:*

- a) *If $\mathcal{E}_1, \mathcal{E}_2$ are subsets of X such that $1_{\mathcal{E}_1}, 1_{\mathcal{E}_2} \in \tilde{\mathcal{D}}_X$, then $1_{\mathcal{E}_1 \cup \mathcal{E}_2} \in \tilde{\mathcal{D}}_X$.*
- b) *If $\mathcal{E}$ is a subset of X such that $1_{\mathcal{E}} \in \tilde{\mathcal{D}}_X$, then $1_{\sigma(\mathcal{E})} \in \tilde{\mathcal{D}}_X$.*
- c) *If $\mathcal{E}$ is a subset of X such that $1_{\mathcal{E}} \in \tilde{\mathcal{D}}_X$, then $1_{\sigma^{-1}(\mathcal{E})} \in \tilde{\mathcal{D}}_X$.*

Proof. a) Let $\mathcal{E}_1, \mathcal{E}_2$ be subsets of X such that $1_{\mathcal{E}_1}, 1_{\mathcal{E}_2} \in \tilde{\mathcal{D}}_X$, then

$$1_{\mathcal{E}_1 \cup \mathcal{E}_2} = 1_{\mathcal{E}_1} + 1_{\mathcal{E}_2} - 1_{\mathcal{E}_1} 1_{\mathcal{E}_2} \in \tilde{\mathcal{D}}_X.$$

b) Let $\mathcal{E}$ be a subset of X such that $1_{\mathcal{E}} \in \tilde{\mathcal{D}}_X$. Set for each $a \in \mathfrak{a}$,

$$\mathcal{E}_a = \{x \in X \mid ax \in \mathcal{E}\}.$$

It is easy to check that

$$\sigma(\mathcal{E}) = \bigcup_{a \in \mathfrak{a}} \mathcal{E}_a.$$

Since $1_{\mathcal{E}_a} = \tilde{\lambda}_a(1_{\mathcal{E}}) \in \tilde{\mathcal{D}}_X$ (cf. Proposition 4.3), it follows from a) that $1_{\sigma(\mathcal{E})} \in \tilde{\mathcal{D}}_X$.

c) Let $\mathcal{E}$ be a subset of X such that $1_{\mathcal{E}} \in \tilde{\mathcal{D}}_X$. It is easy to check that $1_{\sigma^{-1}(\mathcal{E})} = \tilde{\phi}_X(1_{\mathcal{E}})$, so $1_{\sigma^{-1}(\mathcal{E})} \in \tilde{\mathcal{D}}_X$ by Lemma 8.2. $\square$

Proposition 8.4. *If two subshifts X and Y are conjugate, then $\tilde{\mathcal{D}}_X \cong \tilde{\mathcal{D}}_Y$ and $H_X \cong H_Y$.*

Proof. Let $\psi : X \rightarrow Y$ be a conjugacy. Then we can define a $*$ -isomorphism $\Psi : \mathfrak{B}(Y) \rightarrow \mathfrak{B}(X)$ by setting $\Psi(f)(x) = f(\psi(x))$ for every $f \in \mathfrak{B}(Y)$ and every $x \in X$.

Let $\mu \in L(Y)$. Since $C_Y(\mu)$ is clopen and ψ is continuous, $\psi^{-1}(C_Y(\mu))$ is clopen and hence compact. So since $C_X(\nu)$, $\nu \in L(X)$ is a basis for the

topology of $\mathbf{X}$, there exist a finite number of words $\mu_1, \mu_2, \dots, \mu_r \in \mathbf{L}(\mathbf{X})$ such that

$$\psi^{-1}(C_Y(\mu)) = \bigcup_{k=1}^r C_X(\mu_k).$$

Let $\mu, \nu \in \mathbf{L}(\mathbf{Y})$ and let $\mu_1, \dots, \mu_r, \nu_1, \dots, \nu_s \in \mathbf{L}(\mathbf{X})$ such that

$$\psi^{-1}(C_Y(\mu)) = \bigcup_{k=1}^r C_X(\mu_k)$$

and

$$\psi^{-1}(C_Y(\nu)) = \bigcup_{k=1}^s C_X(\nu_k).$$

Since both $\psi \circ \sigma_X = \sigma_Y \circ \psi$, we have that

$$\begin{aligned} \psi^{-1}(C_Y(\mu, \nu)) &= \psi^{-1}(C_Y(\nu)) \cap \sigma_X^{-|\nu|}(\sigma_X^{|\mu|}(\psi^{-1}(C_Y(\mu)))) \\ &= \left(\bigcup_{k=1}^s C_X(\nu_k) \right) \cap \left(\bigcup_{k=1}^r \sigma_X^{-|\nu|}(\sigma_X^{|\mu|}(C_X(\mu_k))) \right), \end{aligned}$$

so it follows from Lemma 8.3 that

$$\Psi(1_{C_Y(\mu, \nu)}) = 1_{\psi^{-1}(C_Y(\mu, \nu))} \in \tilde{\mathcal{D}}_X.$$

Hence $\Psi(\tilde{\mathcal{D}}_Y) \subseteq \tilde{\mathcal{D}}_X$. In the same way we can prove that $\Psi^{-1}(\tilde{\mathcal{D}}_X) \subseteq \tilde{\mathcal{D}}_Y$, so $\Psi(\tilde{\mathcal{D}}_Y) = \tilde{\mathcal{D}}_X$, and thus $\Psi|_{\tilde{\mathcal{D}}_Y} : \tilde{\mathcal{D}}_Y \rightarrow \tilde{\mathcal{D}}_X$ is a $*$ -isomorphism.

Define $T : \mathbf{H}_Y \rightarrow \mathbf{H}_X$ by

$$T(f_a)_{a \in \mathfrak{a}_Y} = \left(\sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \Psi(f_a) \right)_{b \in \mathfrak{a}_X}$$

and $S : \mathbf{H}_X \rightarrow \mathbf{H}_Y$ by

$$S(g_b)_{b \in \mathfrak{a}_X} = \left(\sum_{b \in \mathfrak{a}_X} \tilde{\lambda}_a(\Psi^{-1}(1_{C_X(b)})) \Psi^{-1}(g_b) \right)_{a \in \mathfrak{a}_Y}.$$

Let $a \in \mathfrak{a}_Y$, $b \in \mathfrak{a}_X$ and $x \in \mathbf{Y}$. If $ax \in \mathbf{Y}$ and $(\psi^{-1}(ax))_1 = b$, then

$$\begin{aligned} \psi^{-1}(ax) &= (\psi^{-1}(ax))_1 \sigma(\psi^{-1}(ax)) \\ &= b \psi^{-1}(\sigma(ax)) \\ &= b \psi^{-1}(x) \end{aligned}$$

and thus $b\psi^{-1}(x) \in \mathbf{X}$ and $(\psi(b\psi^{-1}(x)))_1 = a$.

If $b\psi^{-1}(x) \in \mathbf{X}$ and $(\psi(b\psi^{-1}(x)))_1 = a$, then

$$\begin{aligned}\psi(b\psi^{-1}(x)) &= (\psi(b\psi^{-1}(x)))_1 \sigma(\psi(b\psi^{-1}(x))) \\ &= a\psi(\sigma(b\psi^{-1}(x))) \\ &= a\psi(\psi^{-1}(x)) \\ &= ax\end{aligned}$$

and thus $ax \in \mathbf{Y}$ and $(\psi^{-1}(ax))_1 = b$.

Hence $(ax \in \mathbf{Y} \wedge (\psi^{-1}(ax))_1 = b) \Leftrightarrow (b\psi^{-1}(x) \in \mathbf{X} \wedge (\psi(b\psi^{-1}(x)))_1 = a)$.

So

$$\begin{aligned}\Psi^{-1} \left(\tilde{\lambda}_b \left(\Psi \left(1_{C_Y(a)} \right) \right) \right) (x) &= \begin{cases} \Psi \left(1_{C_Y(a)} \right) (b\psi^{-1}(x)) & \text{if } b\psi^{-1}(x) \in \mathbf{X} \\ 0 & \text{if } b\psi^{-1}(x) \notin \mathbf{X} \end{cases} \\ &= \begin{cases} 1 & \text{if } b\psi^{-1}(x) \in \mathbf{X} \wedge (\psi(b\psi^{-1}(x)))_1 = a \\ 0 & \text{else} \end{cases} \\ &= \begin{cases} 1 & \text{if } ax \in \mathbf{Y} \wedge (\psi^{-1}(ax))_1 = b \\ 0 & \text{else} \end{cases} \\ &= \begin{cases} \Psi^{-1} \left(1_{C_X(b)} \right) (ax) & \text{if } ax \in \mathbf{Y} \\ 0 & \text{if } ax \notin \mathbf{Y} \end{cases} \\ &= \tilde{\lambda}_a \left(\Psi^{-1} \left(1_{C_X(b)} \right) \right) (x)\end{aligned}$$

and hence $\Psi^{-1} \left(\tilde{\lambda}_b \left(\Psi \left(1_{C_Y(a)} \right) \right) \right) = \tilde{\lambda}_a \left(\Psi^{-1} \left(1_{C_X(b)} \right) \right)$ for all $a \in \mathfrak{a}_Y$ and $b \in \mathfrak{a}_X$, and thus

$$\begin{aligned}\langle T(f_a)_{a \in \mathfrak{a}_Y}, (g_b)_{b \in \mathfrak{a}_X} \rangle &= \left\langle \left(\sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_b \left(\Psi \left(1_{C_Y(a)} \right) \right) \Psi(f_a) \right)_{b \in \mathfrak{a}_X}, (g_b)_{b \in \mathfrak{a}_X} \right\rangle \\ &= \sum_{b \in \mathfrak{a}_X} \sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_b \left(\Psi \left(1_{C_Y(a)} \right) \right) \Psi(f_a^*) g_b \\ &= \sum_{a \in \mathfrak{a}_Y} \Psi(f_a^*) \sum_{b \in \mathfrak{a}_X} \tilde{\lambda}_b \left(\Psi \left(1_{C_Y(a)} \right) \right) g_b \\ &= \Psi \left(\left\langle (f_a)_{a \in \mathfrak{a}_Y}, \left(\sum_{b \in \mathfrak{a}_X} \tilde{\lambda}_a \left(\Psi^{-1} \left(1_{C_X(b)} \right) \right) \Psi^{-1}(g_b) \right)_{a \in \mathfrak{a}_Y} \right\rangle \right) \\ &= \Psi(\langle (f_a)_{a \in \mathfrak{a}_Y}, S(g_b)_{b \in \mathfrak{a}_X} \rangle)\end{aligned}$$

for all $(f_a)_{a \in \mathfrak{a}_Y} \in \mathbf{H}_Y$ and all $(g_b)_{b \in \mathfrak{a}_X} \in \mathbf{H}_X$.

Let $a \in \mathfrak{a}_Y$, $b \in \mathfrak{a}_X$ and $y \in X$. If $by \in X$ and $(\psi(by))_1 = a$, then

$$\begin{aligned}\psi(by) &= (\psi(by))_1 \sigma(\psi(by)) \\ &= a\psi(\sigma(by)) \\ &= a\psi(y),\end{aligned}$$

and thus $a\psi(y) \in Y$.

So for every $f \in \tilde{\mathcal{D}}_Y$ is

$$\begin{aligned}\tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \Psi(\tilde{\lambda}_a(f))(y) &= \begin{cases} \Psi(1_{C_Y(a)})(by) \Psi(\tilde{\lambda}_a(f))(y) & \text{if } by \in X \\ 0 & \text{if } by \notin X \end{cases} \\ &= \begin{cases} 1_{C_Y(a)}(\psi(by)) \tilde{\lambda}_a(f)(\psi(y)) & \text{if } by \in X \\ 0 & \text{if } by \notin X \end{cases} \\ &= \begin{cases} f(a\psi(y)) & \text{if } by \in X, (\psi(by))_1 = a \text{ and } a\psi(y) \in Y \\ 0 & \text{else} \end{cases} \\ &= \begin{cases} f(\psi(by)) & \text{if } by \in X \text{ and } (\psi(by))_1 = a \\ 0 & \text{else} \end{cases} \\ &= \tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \tilde{\lambda}_b(\Psi(f))(y),\end{aligned}$$

and hence $\tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \Psi(\tilde{\lambda}_a(f)) = \tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \tilde{\lambda}_b(\Psi(f))$ for all $f \in \tilde{\mathcal{D}}_Y$, $a \in \mathfrak{a}_Y$ and $b \in \mathfrak{a}_X$. Thus

$$\begin{aligned}T(\phi'(f)(f_a)_{a \in \mathfrak{a}_Y}) &= T(\tilde{\lambda}_a(f)f_a)_{a \in \mathfrak{a}_Y} \\ &= \left(\sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \Psi(\tilde{\lambda}_a(f)f_a) \right)_{b \in \mathfrak{a}_X} \\ &= \left(\tilde{\lambda}_b(\Psi(f)) \sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \Psi(f_a) \right)_{b \in \mathfrak{a}_X} \\ &= \phi'(\Psi(f))T(f_a)_{a \in \mathfrak{a}_Y}\end{aligned}$$

for all $(f_a)_{a \in \mathfrak{a}_Y} \in H_Y$ and all $f \in \tilde{\mathcal{D}}_Y$.

Since $\Psi^{-1}(\tilde{\lambda}_b(\Psi(1_{C_Y(a)}))) = \tilde{\lambda}_a(\Psi^{-1}(1_{C_X(b)}))$ for all $a \in \mathfrak{a}_Y$ and $b \in \mathfrak{a}_X$, Ψ , Ψ^{-1} and $\tilde{\lambda}_a$ are $*$ -homomorphisms and

$$1_{C_X(b)}1_{C_X(b')} = \begin{cases} 1_{C_X(b)} & \text{if } b = b' \\ 0 & \text{if } b \neq b' \end{cases}$$

for $b, b' \in \mathfrak{a}_X$, we have that

$$\tilde{\lambda}_b(\Psi(1_{C_Y(a)})) \tilde{\lambda}_{b'}(\Psi(1_{C_Y(a)})) = \begin{cases} \tilde{\lambda}_b(\Psi(1_{C_Y(a)})) & \text{if } b = b' \\ 0 & \text{if } b \neq b' \end{cases}$$

for all $a \in \mathfrak{a}_Y$ and all $b, b' \in \mathfrak{a}_X$; and hence

$$\begin{aligned}
TS(g_b)_{b \in \mathfrak{a}_X} &= T \left(\sum_{b \in \mathfrak{a}_X} \tilde{\lambda}_a (\Psi^{-1}(1_{C_X(b)})) \Psi^{-1}(g_b) \right)_{a \in \mathfrak{a}_Y} \\
&= \left(\sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_{b'} (\Psi(1_{C_Y(a)})) \sum_{b \in \mathfrak{a}_X} \Psi(\tilde{\lambda}_a (\Psi^{-1}(1_{C_Y(b)}))) g_b \right)_{b' \in \mathfrak{a}_X} \\
&= \left(\sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_{b'} (\Psi(1_{C_X(a)})) \sum_{b \in \mathfrak{a}_X} \tilde{\lambda}_b (\Psi(1_{C_X(a)})) g_b \right)_{b' \in \mathfrak{a}_X} \\
&= \left(\sum_{a \in \mathfrak{a}_Y} \tilde{\lambda}_b (\Psi(1_{C_Y(a)})) g_b \right)_{b \in \mathfrak{a}_X} \\
&= \left(\tilde{\lambda}_b \left(\Psi \left(\sum_{a \in \mathfrak{a}_Y} 1_{C_Y(a)} \right) \right) g_b \right)_{b \in \mathfrak{a}_X} \\
&= (\tilde{\lambda}_b(1) g_b)_{b \in \mathfrak{a}_X} \\
&= (1_{\sigma(C_X(b))} g_b)_{b \in \mathfrak{a}_X} \\
&= (g_b)_{b \in \mathfrak{a}_X}
\end{aligned}$$

for all $(g_b)_{b \in \mathfrak{a}_X} \in H_X$.

In the same way one can prove that $ST(f_a)_{a \in \mathfrak{a}_Y} = (f_a)_{a \in \mathfrak{a}_Y}$ for all $(f_a)_{a \in \mathfrak{a}_Y} \in H_Y$.

Hence (T, Ψ) is a C^* -correspondence isomorphism and $H_X \cong H_Y$. $\square$

Remark 8.5. With Proposition 8.4 in hand, it is not difficult to prove that if Λ and Γ are two two-sided subshifts such that the one-sided subshift X_Λ of Λ and the one-sided subshift X_Γ of Γ are conjugate, then $\mathcal{O}_\Lambda$ and $\mathcal{O}_\Gamma$ of [5] (cf. Remark 7.5) are isomorphic.

Theorem 8.6. *If two subshifts X and Y are conjugate, then there exists a $*$ -isomorphism ρ from $\mathcal{O}_X$ to $\mathcal{O}_Y$ such that $\gamma_z \circ \rho = \rho \circ \gamma_z$ for every $z \in \mathbb{T}$.*

Proof. It follows from Theorem 8.4 that there exists a C^* -correspondence isomorphism (T, Ψ) from H_X to H_Y . Thus there exists a $*$ -isomorphism $\rho : \mathcal{O}_X \rightarrow \mathcal{O}_Y$ such that $\rho(k_{H_X}(\xi)) = k_{H_Y}(T\xi)$ for every $\xi \in H_X$ and $\rho(k_{\tilde{D}_X}(f)) =$

$k_{\tilde{\mathcal{D}}_Y}(\Psi(f))$ for every $f \in \tilde{\mathcal{D}}_X$. Hence

$$\begin{aligned}
\gamma_z(\rho(k_{H_X}(\xi))) &= \gamma_z(k_{H_Y}(T\xi)) \\
&= zk_{H_Y}(T\xi) \\
&= k_{H_Y}(Tz\xi) \\
&= \rho(k_{H_X}(z\xi)) \\
&= \rho(zk_{H_X}(\xi)) \\
&= \rho(\gamma_z(k_{H_X}(\xi)))
\end{aligned}$$

for every $\xi \in H_X$ and every $z \in \mathbb{T}$, and

$$\begin{aligned}
\gamma_z(\rho(k_{\tilde{\mathcal{D}}_X}(f))) &= \gamma_z(k_{\tilde{\mathcal{D}}_Y}(\Psi(f))) \\
&= zk_{\tilde{\mathcal{D}}_Y}(\Psi(f)) \\
&= k_{\tilde{\mathcal{D}}_Y}(\Psi(zf)) \\
&= \rho(k_{\tilde{\mathcal{D}}_X}(zf)) \\
&= \rho(zk_{\tilde{\mathcal{D}}_X}(f)) \\
&= \rho(\gamma_z(k_{\tilde{\mathcal{D}}_X}(f)))
\end{aligned}$$

for every $f \in \tilde{\mathcal{D}}_X$ and every $z \in \mathbb{T}$. Since $\mathcal{O}_X$ is generated by $k_{H_X}(H_X) \cup k_{\tilde{\mathcal{D}}_X}(\tilde{\mathcal{D}}_X)$, it follows that $\gamma_z \circ \rho = \rho \circ \gamma_z$ for every $z \in \mathbb{T}$. $\square$

Remark 8.7. One can prove that if Λ and Γ are two two-sided subshifts which are flow equivalent (cf. [22], [9], [1] and [12, §13.6]), then $\mathcal{O}_{X_\Lambda}$ and $\mathcal{O}_{X_\Gamma}$, where X_Λ is the one-sided subshift of Λ and X_Γ is the one-sided subshift of Γ , are stably isomorphic. This has been proved in [20] (cf. Remark 7.4) under the assumption of condition (I) and (E) (cf. [19]), but there is an alternative proof, which will appear in [4], which does not require condition (I) and (E).

Notice that if the two two-sided subshifts Λ and Γ are conjugate (as two-sided subshifts, cf. [12, Definition 1.5.9.]), then Λ and Γ are flow equivalent (cf. [12, §13.6]), and so $\mathcal{O}_{X_\Lambda}$ and $\mathcal{O}_{X_\Gamma}$ are stably isomorphic. One can in fact show that if Λ and Γ are conjugate, then there exists a $*$ -isomorphism from Φ from $\mathcal{O}_{X_\Lambda} \otimes \mathcal{K}$ to $\mathcal{O}_{X_\Gamma} \otimes \mathcal{K}$ such that $\Phi(\mathcal{D}_{X_\Lambda} \otimes \mathcal{C}) = \mathcal{D}_{X_\Gamma} \otimes \mathcal{C}$ and such that $\Phi \circ \gamma_\Gamma \circ \Phi^{-1}$ and γ_Λ are exterior equivalent, where $\mathcal{K}$ is the C^* -algebra of all compact operators on a separable infinite-dimensional Hilbert space, $\mathcal{C}$ is a maximal commutative C^* -subalgebra of $\mathcal{K}$, $\mathcal{D}_{X_\Lambda}$ (respectively $\mathcal{D}_{X_\Gamma}$) is the C^* -subalgebra of $\mathcal{O}_\Lambda$ (respectively $\mathcal{O}_\Gamma$) generated by $\{S_\nu S_\mu^* S_\mu S_\nu^*\}_{\mu, \nu \in L(X_\Lambda)}$ (respectively $\{S_\nu S_\mu^* S_\mu S_\nu^*\}_{\mu, \nu \in L(X_\Gamma)}$) (noticed that $\mathcal{D}_{X_\Lambda}$ (respectively $\mathcal{D}_{X_\Gamma}$) is

isomorphic to $\tilde{\mathcal{D}}_{\mathbf{X}_\Lambda}$ (respectively $\tilde{\mathcal{D}}_{\mathbf{X}_\Gamma}$), cf. Remark 7.3) and γ_Λ (respectively γ_Γ) is the gauge action on $\mathcal{O}_{\mathbf{X}_\Lambda}$ (respectively $\mathcal{O}_{\mathbf{X}_\Gamma}$). This has been proved in [19, Corollary 6.2] for two-sided subshifts satisfying the conditions (I) and (E), and will be proved in full generality in [4].

References

- [1] M. Boyle and D. Handelman, *Orbit equivalence, flow equivalence and ordered cohomology*, Israel J. Math. **95** (1996), 169–210.
- [2] T. M. Carlsen, *On C^* -algebras associated with sofic shifts*, J. Operator Theory **49** (2003), 203–212.
- [3] ———, *Operator Algebraic Applications in Symbolic Dynamics*, Ph.D. Thesis, University of Copenhagen, 2004, 189 pp.
- [4] ———, *A faithful representation of the C^* -algebra associated to a shift space* (in preparation.)
- [5] T. M. Carlsen and K. Matsumoto, *Some remarks on the C^* -algebras associated with subshifts*, Math. Scand. **95** (2004), 145–160.
- [6] T. M. Carlsen and S. Silvestrov, *C^* -crossed products and shift spaces*, Preprints in Mathematical Sciences 2005:26, Lund Institute of Technology, 2005, 28 pp.
- [7] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math. **56** (1980), 251–268.
- [8] N. J. Fowler, P. S. Muhly, and I. Raeburn, *Representations of Cuntz-Pimsner algebras*, Indiana Univ. Math. J. **52** (2003), 569–605.
- [9] J. Franks, *Flow equivalence of subshifts of finite type*, Ergodic Theory Dynam. Systems **4** (1984), 53–66.
- [10] Y. Katayama, K. Matsumoto, and Y. Watatani, *Simple C^* -algebras arising from β -expansion of real numbers*, Ergodic Theory Dynam. Systems **18** (1998), 937–962.
- [11] T. Katsura, *A construction of C^* -algebras from C^* -correspondences*, Advances in Quantum Dynamics, 173–182, Contemp. Math, 335, Amer. Math. Soc., Providence, RI, 2003.
- [12] D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press, Cambridge, 1995, ISBN 0-521-55124-2, 0-521-55900-6.
- [13] K. Matsumoto, *On C^* -algebras associated with subshifts*, Internat. J. Math. **8** (1997), 357–374.
- [14] ———, *K-theory for C^* -algebras associated with subshifts*, Math. Scand. **82** (1998), 237–255.
- [15] ———, *A simple C^* -algebra arising from a certain subshift*, J. Operator Theory **42** (1999), 351–370.
- [16] ———, *Dimension groups for subshifts and simplicity of the associated C^* -algebras*, J. Math. Soc. Japan **51** (1999), 679–698.
- [17] ———, *Relations among generators of C^* -algebras associated with subshifts*, Internat. J. Math. **10** (1999), 385–405.
- [18] ———, *On automorphisms of C^* -algebras associated with subshifts*, J. Operator Theory **44** (2000), 91–112.
- [19] ———, *Stabilized C^* -algebras constructed from symbolic dynamical systems*, Ergodic Theory Dynam. Systems **20** (2000), 821–841.
- [20] ———, *Bowen-Franks groups for subshifts and Ext-groups for C^* -algebras*, K-Theory **23** (2001), 67–104.

- [21] K. Matsumoto, Y. Watatani, and M. Yoshida, *KMS states for gauge actions on C^* -algebras associated with subshifts*, Math. Z. **228** (1998), 489–509.
- [22] B. Parry and D. Sullivan, *A topological invariant of flows on 1-dimensional spaces*, Topology **14** (1975), 297–299.
- [23] M. V. Pimsner, *A class of C^* -algebras generalizing both Cuntz-Krieger algebras and crossed products by $\mathbf{Z}$* , Free Probability Theory (Waterloo, ON, 1995), Fields Inst. Commun., vol. 12, Amer. Math. Soc., Providence, RI, 1997, pp. 189–212.
- [24] J. Schweizer, *Dilations of C^* -correspondences and the simplicity of Cuntz-Pimsner algebras*, J. Funct. Anal. **180** (2001), 404–425.